

Influence of Extratropical Thermal Forcings on the Asian Monsoons

Stefanie TALENTO^{1,2}, and Jürg LUTERBACHER¹

¹ *Justus Liebig University of Giessen, Giessen, Germany*

² *University of the Republic, Montevideo, Uruguay*

The capability of an extratropical thermal forcing to affect the remote tropical climate has been suggested in studies analysing paleoclimatic data, twentieth century observations and numerical simulations. The Intertropical Convergence Zone (ITCZ) has been the most studied feature and the general picture emerging from these studies is that the ITCZ tends to shift toward the warmer hemisphere. However, there is evidence supporting the idea that other features of the tropical climate might be affected by such an extratropical influence. In particular, the Asian Summer Monsoons (ASM) stand out as some of such possibly affected systems.

This study investigates, through numerical simulations, the response of the ASM to a series of extratropical thermal forcings. Three sets of experiments are performed, varying the extratropical forcing. In the first set the forcing consists of warming of the Northern Hemisphere (NH) extratropics (poleward of 40°) and cooling of the Southern Hemisphere extratropics, with zero global average. In the second and third experiments, the former forcing is divided into its northern and southern components to assess their relative roles in affecting the Asian summer climate. In all the cases realistic boundary surface conditions are used.

The numerical configuration consists of an Atmospheric General Circulation Model (ICTP-AGCM) coupled to either a Slab Ocean Model or a combination of a Slab and a Reduced Gravity Ocean Model. The relative roles of the atmosphere, tropical Sea Surface Temperatures (SSTs), continental surface temperatures and tropical ocean dynamics are investigated in a series of experiments designed to separate these influences.

We find that the imposed extratropical forcings generate a response in the ASM. This response is strongest over central China and South-eastern Asia, while over India the forced behaviour is weak and limited in spatial extension. When a NH extratropical warming is imposed we find that a precipitation dipole-like pattern is generated, with wet (dry) conditions over central China (South-eastern Asia). This precipitation dipole structure is exacerbated when the tropical SSTs are not allowed to react to the remote extratropical forcing, suggesting that the atmospheric dynamics behaviour is weakened by the surface oceanic thermodynamic response. The relative roles of the land surface temperatures, the tropical ocean dynamics and the hemispheric components of the forcing are also investigated.

This research is carried out within a project funded under the 2015 joint JPI Climate-Belmont Forum call (INTEGRATE, An integrated data-model study of interactions between tropical monsoons and extra-tropical climate variability and extremes).

Key words: Asian Monsoons, extratropical forcing, teleconnection, slab